

GRAM POSITIVE

Antimicrobial susceptibility results

1/1/2017 - 12/31/2017

	<i>Enterococcus faecalis</i>	<i>Enterococcus faecium</i>	<i>Staphylococcus aureus</i>	MRSA	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus lugdunensis</i>	Group B <i>Streptococcus</i>	<i>Streptococcus pneumoniae</i>
Max strains tested	569	56	1,542	418	98	69	83	61
% Susceptible								
Amoxicillin/clavulanate	-	-	-	0	-	-	-	93
Ampicillin	99	39	-	0	-	-	-	-
Ciprofloxacin	72*	27*	76	38	54	97	-	-
Clindamycin	-	-	74	67	59	93	52	92
Erythromycin	-	-	52**	12**	30**	91**	-	56
Linezolid	100	100	100	99	100	-	-	-
Oxacillin/Nafcillin	-	-	73	0	33	91	-	-
Tetracycline	31	-	95***	92***	86***	97***	-	79
Trimethoprim/sulfa	-	-	96	95	44	100	-	-
Vancomycin	100	57	100	100	99	100	-	100
Penicillin (CSF)	-	-	-	-	-	-	-	72
Penicillin (Non-CSF)	-	-	-	-	-	-	-	98
Ceftriaxone (CSF)	-	-	-	-	-	-	-	85
Ceftriaxone (Non-CSF)	-	-	-	-	-	-	-	100
Cefazolin	-	-	73	0	33	89	-	-
For urinary tract infections only								
Nitrofurantoin	97	16	97	95	97	99	-	-

\*Should only be used to treat urinary tract infections caused by *Enterococcus* species.

\*\*Susceptibility to erythromycin can be used to infer susceptibility to azithromycin.

\*\*\*Susceptibility to tetracycline can be used to infer susceptibility to doxycycline.